

UNIT – I
RELATIONS & FUNCTIONS

Multiple Choice Questions(1 Mark)

- 1 Relation $R = \{(x, y) : x \leq y, x, y \in \mathbb{Z}\}$ is
 - (a)Reflexive and symmetric relation
 - (b)Symmetric and transitive relation
 - (c)Equivalence relation
 - (d)Reflexive and transitive relation
- 2 Which of the following relations defined on set $A = \{1, 2, 3\}$ is reflexive but neither symmetric nor transitive :
 - (a) $R = \{(1, 1), (2, 2), (3, 3)\}$
 - (b) $R = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3)\}$
 - (c) $R = \{(1, 2), (1, 3), (2, 3), (3, 1), (2, 1)\}$
 - (d) $R = \{(1, 2), (2, 3), (1, 3), (2, 1)\}$
- 3 Function defined by $f : R \rightarrow R, f(x) = x^3$ is :
 - (a)only one-one
 - (b)only onto
 - (c)one-one and onto
 - (d)neither one-one nor onto
- 4 Function defined by $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$ is :
 - (a)only one-one
 - (b)only onto
 - (c)one-one and onto
 - (d)neither one-one nor onto
- 5 Relation $R = \{(x, x), (y, y), (x, y), (y, x)\}$ defined on the set $A = \{x, y\}$ is :
 - (a)Only Reflexive relation
 - (b)Only Symmetric relation
 - (c)Only Transitive relation
 - (d)Equivalence relation
- 6 Relation $R = \{(x, y) : x < y, x, y \in \mathbb{Z}\}$ is
 - (a)Only Reflexive relation
 - (b)Only Symmetric relation
 - (c)Only Transitive relation
 - (d)Equivalence relation
- 7 Relation $R = \{(x, y) : x < y^2 \text{ where } x, y \in \mathbb{R}\}$ is
 - (a)Reflexive but not symmetric
 - (b)Symmetric and transitive but not Reflexive
 - (c)Reflexive and Symmetric
 - (d)Neither reflexive nor symmetric nor transitive
- 8 If $A = \{1, 4, 9, 16, 25, \dots\}$ then function defined by $f : \mathbb{Z} \rightarrow A, f(x) = x^2$ is
 - (a)only one-one
 - (b)only onto
 - (c)function is not defined
 - (d)neither one-one nor onto
- 9 If $A = \{0, 1, 4, 9, 16, 25, \dots\}$ then function defined by $f : \mathbb{N} \rightarrow A, f(x) = x^2$ is
 - (a)one-one but not onto
 - (b)onto but not one-one
 - (c)one-one and onto
 - (d)neither one-one nor onto
- 10 Function $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = \frac{3-7x}{2}$ is:
 - (a)one-one but not onto
 - (b)onto but not one-one
 - (c)one-one and onto
 - (d)neither one-one nor onto
- 11 If $A = \{0, 1, 4, 9, 16, 25, \dots\}$ then function defined by $f : \mathbb{Z} \rightarrow A, f(x) = x^2$ is
 - (a)one-one but not onto
 - (b)onto but not one-one
 - (c)one-one and onto
 - (d)neither one-one nor onto
- 12 If $A = \{1, 4, 9, 16, 25, \dots\}$ then function defined by $f : \mathbb{N} \rightarrow A, f(x) = x^2$ is
 - (a)only one-one
 - (b)only onto
 - (c)one-one and onto
 - (d)neither one-one nor onto
- 13 Function $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = \frac{1}{x}$ is:
 - (a)one-one but not onto
 - (b)onto but not one-one
 - (c)function is not defined
 - (d)neither one-one nor onto
- 14 Relation $R = \{(x, y) : x \leq y^3 \text{ where } x, y \in \mathbb{R}\}$ is
 - (a)Reflexive but not symmetric
 - (b)Symmetric and transitive but not Reflexive
 - (c)Reflexive and Symmetric
 - (d)Neither reflexive nor symmetric nor transitive
- 15 Function defined by $f : \mathbb{Z} \rightarrow \mathbb{W}, f(x) = x^2$ is
 - (a)one-one but not onto
 - (b)onto but not one-one
 - (c)one-one and onto
 - (d)neither one-one nor onto

Fill in the blanks(1 Mark)

2

- 1) Identity relation is also _____
- 2) If R is a relation defined on the set A then R is a subset of _____
- 3) If function f is defined as $f: A \rightarrow f(A)$ then f is a _____ function.
- 4) If R is a relation from A to B then R is _____ of $A \times B$.
- 5) The relation R (defined on the set A) is called _____ if $(x, x) \in R \ \forall x \in A$
- 6) The relation R (defined on the set A) is called _____ if $(x, y) \in R \Rightarrow (y, x) \in R$
 $\forall x, y \in A$
- 7) The relation R (defined on the set A) is called _____ if $(x, y), (y, z) \in R \Rightarrow (x, z) \in R$
 $\forall x, y, z \in A$
- 8) If for the numbers x_1, x_2 in the domain of the function f we have $f(x_1) = f(x_2)$ implies $x_1 = x_2$ then function is _____

4 Marks Questions

1. Check reflexivity, symmetry and transitivity for the following relations :
 - (i) $R = \{(x, y) : x - y \text{ is an integer}\}$ (defined on the set of integers \mathbb{Z})
 - (ii) $R = \{(x, y) : |x - y| \text{ is an integer}\}$ (defined on the set of integers \mathbb{Z})
 - (iii) $R = \{(x, y) : x - y \text{ is divisible by 3}\}$ (defined on the set of integers \mathbb{Z})
 - (iv) $R = \{(x, y) : |x - y| \text{ is divisible by 6}\}$ (defined on the set of integers \mathbb{Z})
 - (v) $R = \{(x, y) : x \leq y^2 \text{ where } x, y \in \mathbb{R}\}$
 - (vi) $R = \{(x, y) : x \leq y^3 \text{ where } x, y \in \mathbb{R}\}$
 - (vii) $R = \{ (l_1, l_2) : \text{line } l_1 \text{ is parallel to the line } l_2 \}$ (Defined on the set of all lines L)
 - (viii) $R = \{ (l_1, l_2) : \text{line } l_1 \text{ is perpendicular to the line } l_2 \}$ (Defined on the set of all lines L in a plane)
2. For the following functions $f : R \rightarrow R$:
 - (i) $f(x) = \frac{3x+5}{2}$
 - (ii) $f(x) = \frac{2x-7}{4}$
 - (iii) $f(x) = \frac{3-2x}{4}$
 - (iv) $f(x) = \frac{4-3x}{5}$
 - (v) $f(x) = \frac{6-5x}{7}$
 - (vi) $f(x) = \frac{5x+7}{6}$

show that these functions are one-one and onto.

UNIT – I

Inverse Trigonometric Functions

Multiple Choice Questions(1 Marks Questions)

1 $\cos^{-1} \left(\cos \frac{2\pi}{3} \right)$ is equal to :
 (a) $\frac{\pi}{5}$ (b) $\frac{2\pi}{3}$ (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{3}$

2 $\sin^{-1} \left(\frac{1}{2} \right)$ is equal to :
 (a) 0 (b) $\frac{\pi}{6}$ (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{3}$

3 $\cos^{-1}(0)$ is equal to :
 (a) 0 (b) $\frac{\pi}{6}$ (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{3}$

4 $\tan^{-1}(1)$ is equal to :
 (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{6}$ (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{3}$

5 If $y = \sin^{-1}(x)$ then x belongs to the interval :
 (a) $(0, \pi)$ (b) $(-1, 1)$ (c) $[-1, 1]$ (d) $[0, \pi]$

6 $\sin^{-1} \left(\sin \frac{\pi}{3} \right)$ is equal to :
 (a) $\frac{\pi}{5}$ (b) $\frac{2\pi}{3}$ (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{3}$

7 If $\cos^{-1} x = y$ then x belongs to
 (a) $(0, 1)$ (b) $(-1, 1)$ (c) $[-1, 1]$ (d) $[0, 1]$

8 Principal value of $\cos^{-1} \left(\cos \left(\frac{7\pi}{4} \right) \right)$ is
 (a) 0 (b) $\frac{7\pi}{4}$ (c) $\frac{\pi}{4}$ (d) $\frac{\pi}{6}$

9 Range of function \sec^{-1} is :
 (a) $[0, \pi] - \left\{ \frac{\pi}{2} \right\}$ (b) $(0, \pi)$ (c) $\left(-\frac{\pi}{2}, \frac{\pi}{2} \right) - \{0\}$ (d) $[0, \pi]$

10 Domain of function $\operatorname{cosec}^{-1}$ is :
 (a) $[-1, 1]$ (b) $\mathbb{R} - (-1, 1)$ (c) \mathbb{R} (d) $(-1, 1)$

11 Domain of the function \tan^{-1} is :
 (a) $\left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$ (b) $\mathbb{R} - (-1, 1)$ (c) \mathbb{R} (d) $(-1, 1)$

12 If $\tan^{-1} x = y$, then y belongs to the interval :
 (a) $\left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$ (b) $\mathbb{R} - (-1, 1)$ (c) \mathbb{R} (d) $(-1, 1)$

4 Marks Questions

1. Find the values of :

- $5 \sec^{-1}(\sqrt{2}) + 8 \tan^{-1}(1) - 3 \sin^{-1}\left(\sin\left(\frac{3\pi}{4}\right)\right)$
- $2 \operatorname{cosec}^{-1}(1) - 5 \sec^{-1}\left(\frac{2}{\sqrt{3}}\right) + \sin^{-1}\left(\frac{1}{2}\right) - 4 \cot^{-1}(\sqrt{3})$
- $3 \operatorname{cosec}^{-1}(1) + \sec^{-1}(2) - 5 \sin^{-1}\left(\frac{\sqrt{3}}{2}\right) + 7 \cot^{-1}\left(\frac{1}{\sqrt{3}}\right)$

2. Prove that : (i) $\cos^{-1}\frac{4}{5} + \cos^{-1}\frac{12}{13} = \cos^{-1}\frac{33}{65}$ (ii) $\sin^{-1}\frac{8}{17} + \sin^{-1}\frac{3}{5} = \tan^{-1}\frac{77}{36}$

UNIT – II (Algebra)
MATRICES & DETERMINANTS
Multiple Choice Questions(1 Mark)

1 If order of matrix A is 2×3 and order of matrix B is 3×5 then order of matrix $B'A'$ is :
 (a) 5×2 (b) 2×5 (c) 5×3 (d) 3×2

2 If $\begin{vmatrix} x & 1 \\ 1 & x \end{vmatrix} = \begin{vmatrix} 2 & 0 \\ 8 & 4 \end{vmatrix}$ then value of x is :
 (a) 3 (b) 2 (c) 4 (d) 8

3 If $\begin{bmatrix} 2x+y & 0 \\ 5 & x \end{bmatrix} = \begin{bmatrix} 5 & 0 \\ 5 & 3 \end{bmatrix}$, then y is equal to:-
 (a) 1 (b) 3 (c) 2 (d) -1

4 If $A + B = C$ where B and C are matrices of order 5×5 then order of A is :-
 (a) 5×5 (b) 5×3 (c) 3×5 (d) 3×3

5 If $A B = C$ where B and C are matrices of order 2×5 and 5×5 respectively then order of A is :-
 (a) 5×5 (b) 5×2 (c) 2×5 (d) 2×2

6 If order of matrix A is 2×3 and order of matrix B is 3×5 then order of matrix AB is :
 (a) 5×2 (b) 2×5 (c) 5×3 (d) 3×2

7 If order of matrix A is 4×3 and order of matrix B is 3×5 then order of matrix AB is :
 (a) 5×4 (b) 4×5 (c) 5×3 (d) 3×4

8 If A is a square matrix of order 4×4 and $|A| = 3$ then $|Adj.(A)|$ is
 (a) 27 (b) 81 (c) 9 (d) 3

9 If $A = \begin{bmatrix} 2 & 5 \\ 1 & -2 \end{bmatrix}$ then $|A|$ is
 (a) -9 (b) 9 (c) 1 (d) -1

10 If A is a matrix of order of 3×3 and $|A| = 3$ then $|Adj(A)|$ is
 (a) 81 (b) 9 (c) 27 (d) 3

Fill-ups(1 Mark)

- 1) If $A = [a_{ij}]_{2 \times 3}$ such that $a_{ij} = i + j$ then $a_{11} = \underline{\hspace{2cm}}$
- 2) If $|A| = 5$ where A is a matrix of order 3×3 then $|adj.(A)| = \underline{\hspace{2cm}}$
- 3) If matrix $A = \begin{bmatrix} 2 & 3 \\ 1 & 5 \end{bmatrix}$ then $|A| = \underline{\hspace{2cm}}$
- 4) If order of matrix A is 3×4 then order of $A' = \underline{\hspace{2cm}}$
- 5) If for a matrix $, A' = A$ holds then A is $\underline{\hspace{2cm}}$ matrix.
- 6) If for a matrix $, A' = -A$ holds then A is $\underline{\hspace{2cm}}$ matrix.
- 7) If for any two matrices A and B , $AB = BA = I$ then these matrices are $\underline{\hspace{2cm}}$ of each other.
- 8) $\underline{\hspace{2cm}}$ matrix is symmetric as well as skew-symmetric.
- 9) If order of matrix A is 3×4 and order of matrix B is 4×7 then order of AB is $\underline{\hspace{2cm}}$.
- 10) If order of matrix A is 4×5 then number of elements in A are $\underline{\hspace{2cm}}$

2 Marks Questions

5

1. If $A = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix}$, then verify $A^2 - 7A - 2I = 0$.
2. If $A = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$ show that $A^2 - 5A - 14I = 0$.
3. If $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ and $f(x) = x^2 - 4x + 1$ then find $f(A)$.
4. If $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $f(x) = x^2 - 2x - 3$ then find $f(A)$.
5. If $A = \begin{bmatrix} 1 & 0 \\ -1 & 7 \end{bmatrix}$ and $A^2 - 8A = kI$ then find k .
6. If $A = \begin{bmatrix} 0 & 3 \\ -7 & 5 \end{bmatrix}$ then find k so that $kA^2 = 5A - 21I$.
7. If $X = \begin{bmatrix} 3 & 4 \\ 2 & -1 \end{bmatrix}$ and $2X - Y = \begin{bmatrix} 5 & 10 \\ 3 & -5 \end{bmatrix}$ then find the matrix Y .
8. If $X - 2Y = \begin{bmatrix} 5 & 1 \\ 2 & 0 \end{bmatrix}$ and $2X - Y = \begin{bmatrix} 4 & 9 \\ 1 & -3 \end{bmatrix}$ then find the matrices X and Y .

9. Verify $(AB)' = B'A'$ for the following matrices :

- $A = \begin{bmatrix} 1 \\ 3 \\ 6 \end{bmatrix}$, $B = [2 \quad 4 \quad 5]$
- $A = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix}$, $B = [-2 \quad -1 \quad -4]$
- $A = \begin{bmatrix} 2 & 3 \\ 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix}$
- $A = \begin{bmatrix} 4 & -2 \\ -1 & 3 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 7 \\ 5 & 0 \end{bmatrix}$
- $A = \begin{bmatrix} -1 & 3 & 0 \\ -7 & 2 & 8 \end{bmatrix}$, $B = \begin{bmatrix} -5 & 0 \\ 0 & 3 \\ 1 & -8 \end{bmatrix}$
- $A = \begin{bmatrix} 2 & 1 & 3 \\ 4 & 1 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & -1 \\ 0 & 2 \\ 5 & 0 \end{bmatrix}$

10. Using determinants, show that following points are collinear :

- (11, 7), (5, 5) and (-1, 3)
- (3, 8), (-4, 2) and (10, 14)
- (-2, 5), (-6, -7) and (-5, -4)

11. Find the value of x if (3, -2), (x , 2) and (8, 8) are collinear points.

12. Using determinants, find the value of k if the area of the triangle formed by the points (-3, 6), (-4, 4) and (k , -2) is 12 sq. units.

13. If the area of triangle is 35 sq. units with vertices (2, -6), (5, 4) and (k , 4) then find the value of k .

14. Find the equation of the line passing from (3, 2) and (-4, -7) using determinants.

6/4 Marks Questions

6

1. Solve the following system of linear equations by matrix method :

(i) $x - y + 2z = 7, 3x + 4y - 5z = -5, 2x - y + 3z = 12$

(ii) $x + y + z = 6, y + 3z = 11, x - 2y + z = 0$

(iii) $3x + y + z = 10, 2x - y - z = 0, x - y + 2z = 1$

(iv) $2x + 3y + 3z = 5, x - 2y + z = -4, 3x - y - 2z = 3$

(v) $\frac{2}{x} + \frac{3}{y} + \frac{3}{z} = 5, \frac{1}{x} - \frac{2}{y} + \frac{1}{z} = -4, \frac{3}{x} - \frac{1}{y} - \frac{2}{z} = 3$

(vi) $x - y + 2z = 2, 3x + 4y - 5z = 2, 2x - y + 3z = 4$

(vii) $x + y - z = 3, 2x + 3y + z = 10, 3x - y - 7z = 1$

(viii) $x + y + z = 3, 5x - y - z = 3, 3x + 2y - 4z = 1$

2. Express the following matrices as a sum of a symmetric matrix and a skew-symmetric matrix :

(i) $\begin{bmatrix} 2 & 0 & 3 \\ -1 & 4 & 8 \\ 7 & 2 & 9 \end{bmatrix}$

(ii) $\begin{bmatrix} 3 & 6 & 2 \\ 0 & 7 & 8 \\ 5 & 1 & 9 \end{bmatrix}$

(iii). $\begin{bmatrix} 5 & 1 & 2 \\ -2 & 3 & 0 \\ 6 & 3 & 7 \end{bmatrix}$

(iv) $\begin{bmatrix} 2 & 5 & 8 \\ -3 & 6 & 0 \\ 5 & 2 & 1 \end{bmatrix}$

(v). $\begin{bmatrix} 4 & 5 \\ 7 & 9 \end{bmatrix}$

(vi) $\begin{bmatrix} 7 & -3 \\ 4 & 5 \end{bmatrix}$

3. Find the adjoint matrix of the following matrices :

(i) $\begin{bmatrix} 5 & 5 & 2 \\ -1 & 4 & 4 \\ 7 & 0 & 9 \end{bmatrix}$

(ii) $\begin{bmatrix} 2 & 5 & 9 \\ 3 & 7 & 8 \\ 5 & 0 & 9 \end{bmatrix}$

(iii). $\begin{bmatrix} 5 & 1 & 2 \\ -2 & 3 & 0 \\ 6 & 3 & 7 \end{bmatrix}$

(iv) $\begin{bmatrix} 2 & 5 & 8 \\ -3 & 6 & 0 \\ 5 & 2 & 1 \end{bmatrix}$

4. Show that $A = \begin{bmatrix} 2 & -3 \\ 3 & 4 \end{bmatrix}$ and $f(x) = x^2 - 6x + 17$ then show that $f(A) = O$. Using this result find

the matrix A^{-1} .

5. If $A = \begin{bmatrix} 3 & 2 & 0 \\ 1 & 4 & 0 \\ 0 & 0 & 5 \end{bmatrix}$ and $f(x) = x^2 - 7x + 10$ then show that $f(A) = O$. Using this result find

the matrix A^{-1} .

6. Ajay, Sameer and Meenal have Rs.20/- each and some toy guns, chess sets and rubic cubes in their shops. In a week, Ajay sold 3 toy guns and a rubic cube but he bought 2 chess sets for his shop and he has Rs.35/- now. In same duration, Sameer sold 2 chess sets and 2 rubic cubes but he bought a toy gun for his shop and he has Rs. 95/- now. Similarly, Meenal sold 2 toy guns and a chess set but she bought 3 rubic cubes for her shop and she has Rs.15/- now. Find the cost of a toy gun, a chess set and a rubic cube by the help of matrices.
7. If present ages of Rekha and Sanju is subtracted from the present age of Renu then we get 2. Two years ago, if we subtract age of Rekha from the sum of the ages of Renu and Sanju then we get 8. Four years ago, age of Renu was four times the sum of the ages of Rekha and Sanju. Represent the situation algebraically and find the present age of Renu, Rekha and Sanju using matrices.
8. Ajay bought 3 pencils, 2 scales and an eraser for Rs.25/- . Ravinder bought a pencil, 3 scales and 2 erasers for Rs.22/- . Razzaak bought 4 pencils and 3 erasers for Rs. 32/- . Find the costs of a pencil, a scale and an eraser by using matrix method.
9. Two points are moving along straight paths. Point 1 passes through the points (0,5) and (2,3). Point 2 passes through the points (1,0) and (5,4). Find the loci of paths of both the points. Also find the coordinates where both points coincides using matrix method.
10. The sum of three numbers is 2. If twice the second number is added to the sum of first and third, the sum is 1. By adding second and third number to five times the first number , we get 6. Find the three numbers by using matrices.

UNIT – III
Continuity & Differentiability
Multiple Choice Questions (1 Marks)

1 If $f(x) = \begin{cases} kx + 1, & x \leq 5 \\ 3x - 5, & x > 5 \end{cases}$ is continuous then value of k is :
 (a) $\frac{9}{5}$ (b) $\frac{5}{9}$ (c) $\frac{5}{3}$ (d) $\frac{3}{5}$

2 If $f(x) = \begin{cases} kx^2, & x \leq 2 \\ 3, & x > 2 \end{cases}$ is continuous then value of k is :
 (a) $\frac{2}{3}$ (b) $\frac{4}{3}$ (c) $\frac{3}{2}$ (d) $\frac{3}{4}$

3 If $f(x) = \begin{cases} mx - 1, & x \leq 5 \\ 3x - 5, & x > 5 \end{cases}$ is continuous then value of m is :
 (a) $\frac{11}{5}$ (b) $\frac{5}{11}$ (c) $\frac{5}{3}$ (d) $\frac{3}{5}$

4 If $f(x) = \begin{cases} mx^2, & x \leq 5 \\ 6x - 5, & x > 5 \end{cases}$ is continuous then value of m is :
 (a)- 1 (b) 4 (c) 3 (d) 1

5 If $f(x) = \begin{cases} \frac{\sin 2x}{3x}, & x \neq 0 \\ m - 1, & x = 0 \end{cases}$ is continuous then value of m is :
 (a) $2/3$ (b) $3/2$ (c) $3/5$ (d) $5/3$

6 If $f(x) = \begin{cases} kx + 1, & x \leq 5 \\ 3x + 5, & x > 5 \end{cases}$ is continuous then value of k is :
 (a) $\frac{19}{5}$ (b) $\frac{5}{9}$ (c) $\frac{5}{3}$ (d) $\frac{3}{5}$

7 If $f(x) = \begin{cases} kx - 1, & x \leq 5 \\ 3x + 5, & x > 5 \end{cases}$ is continuous then value of k is :
 (a) $\frac{21}{5}$ (b) $\frac{5}{19}$ (c) $\frac{5}{21}$ (d) $\frac{19}{5}$

8 If $f(x) = \begin{cases} \frac{\sin 7x}{3x}, & x \neq 0 \\ m, & x = 0 \end{cases}$ is continuous at $x = 0$ then value of m is
 (a) $\frac{3}{7}$ (b) $\frac{4}{7}$ (c) $\frac{7}{4}$ (d) $\frac{7}{3}$

9 If $y = \log [x + \sqrt{x^2 + 1}]$ then $\frac{dy}{dx}$ is
 (a) $\sqrt{x^2 + 1}$ (b) $\frac{1}{\sqrt{x^2 + 1}}$ (c) $\frac{x}{\sqrt{x^2 + 1}}$ (d) $\frac{1}{x + \sqrt{x^2 + 1}}$

10 If $f(x) = \begin{cases} \frac{x^3 - 8}{x - 2}, & x \neq 2 \\ k, & x = 2 \end{cases}$ is continuous at $x = 2$ then value of k is
 (a) 8 (b) 2 (c) 6 (d) 12

11 $\frac{d}{dx} \{\tan^{-1}(e^x)\}$ is equal to :
 (a) $e^x \tan^{-1} e^x$ (b) $\frac{e^x}{1+e^{2x}}$ (c) 0 (d) $e^x \sec^{-1} x$

12 If $y = \sin x$ then at $x = \frac{\pi}{2}$, y_2 is equal to :
 (a) -1 (b) 1 (c) 0 (d) $\frac{1}{2}$

13 If $x = 2at$, $y = at^2$ then $\frac{dy}{dx}$ is equal to:
 (a) 2 (b) $2a$ (c) $2at$ (d) t

14 If $y = \cos^{-1}(e^x)$ then $\frac{dy}{dx}$ is equal to:
 (a) $e^x \sin^{-1}(e^x)$ (b) $e^x \cos^{-1}(e^x)$ (c) $\frac{-e^x}{\sqrt{1-e^{2x}}}$ (d) $\frac{e^x}{\sqrt{1-e^{2x}}}$

15 If $y = \sin^{-1}(e^x)$ then $\frac{dy}{dx}$ is equal to:
 (a) $e^x \sin^{-1}(e^x)$ (b) $e^x \cos^{-1}(e^x)$ (c) $\frac{-e^x}{\sqrt{1-e^{2x}}}$ (d) $\frac{e^x}{\sqrt{1-e^{2x}}}$

16 $\frac{d}{dx} \{\cot^{-1}(e^x)\}$ is equal to :
 (a) $e^x \tan^{-1} e^x$ (b) $\frac{e^x}{1+e^{2x}}$ (c) $\frac{-e^x}{1+e^{2x}}$ (d) $e^x \sec^{-1} x$

17 If $y = x^2$ then $y_1(5)$ is equal to :
 (a)10 (b)25 (c)32 (d) 0

18 If $y = \log(\sin x)$ then at $x = \frac{\pi}{4}$, $\frac{dy}{dx}$ is
 (a)0 (b)-1 (c)1 (d) $\sqrt{2}$

19 If $y = e^{\log x}$ then $\frac{dy}{dx}$ is
 (a) $\log x - x$ (b) $xe^{\log x}$ (c)1 (d) $e^{\log x} \log x$

20 If $y = \log(\sec x)$ then $\frac{dy}{dx}$ at $x = \frac{\pi}{4}$ is
 (a)1 (b)-1 (c)0 (d)10

2/3 Marks Questions

1. Find the relation between a and b if $f(x) = \begin{cases} ax + 1, & \text{if } x \leq 3 \\ bx + 3, & \text{if } x > 3 \end{cases}$ is a continuous at $x = 3$.

2. Find the values of a and b if the following function is continuous :

$$f(x) = \begin{cases} 5, & \text{if } x \leq 2 \\ ax + b, & \text{if } 2 < x < 10 \\ 21, & \text{if } x \geq 10 \end{cases}$$

3. Find $\frac{dy}{dx}$ for the following parametric functions :

- (i) $x = a \cos^2 \theta, y = b \sin^2 \theta$
- (ii) $x = a(\theta - \sin \theta), y = b(1 + \cos \theta)$
- (iii) $x = a(\theta + \sin \theta), y = b(1 + \cos \theta)$
- (iv) $x^2 + y^2 + 2xy = 23$
- (v) $y = \sin^{-1} \left(\frac{2x}{1+x^2} \right)$
- (vi) $y = \tan^{-1} \left(\frac{2x}{1-x^2} \right)$
- (vii) $x^3 + 3x^2y + 3xy^2 + y^3 = 81$

4. Determine the value of constants a & b so that function f defined below is continuous everywhere:

- (i) $f(x) = \begin{cases} x + 2, & x \leq 2 \\ ax + b, & 2 < x < 5 \\ 3x - 2, & x \geq 5 \end{cases}$
- (ii) $f(x) = \begin{cases} ax^2 + b, & x > 2 \\ 2, & x = 2 \\ 2ax - b, & x < 2 \end{cases}$
- (iii) $f(x) = \begin{cases} 3ax + b, & x > 1 \\ 11, & x = 1 \\ 5ax - 2b, & x < 1 \end{cases}$

5. If $y = \sqrt{x + \sqrt{x + \sqrt{x + \dots \dots \dots \text{to } \infty}}}$ then prove that $(2y - 1) \frac{dy}{dx} = 1$.

6. If $y = \sqrt{3^x + \sqrt{3^x + \sqrt{3^x + \dots \dots \dots \infty}}}$ then prove that $(2y - 1) \frac{dy}{dx} = 3^x \log 3$.

7. If $y = \sqrt{\tan x + \sqrt{\tan x + \sqrt{\tan x + \dots \dots \dots \infty}}}$ then prove that $(2y - 1) \frac{dy}{dx} = \sec^2 x$.

4 Marks Questions

1. If $x = 2 \cos \theta - \cos 2\theta$, $y = 2 \sin \theta - \sin 2\theta$ then find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{2}$.

2. If $x = \frac{1-t^2}{1+t^2}$, $y = \frac{2t}{1+t^2}$ then prove that $\frac{dy}{dx} + \frac{x}{y} = 0$.

3. Differentiate the following w.r.t. :

(i) $x^{\sin x} + (\sin x)^x$ (ii) $x^{\log x} + (\log x)^x$ (iii) $x^{\tan x} + (\tan x)^x$

(iv) $x^{\cos x} + (\sin x)^{\tan x}$ (v) $x^x + (\sin x)^x$ (vi) $x^{\sin^{-1} x} + (\sin^{-1} x)^x$

(vii) $\left(x + \frac{1}{x}\right)^x + x^{\left(1+\frac{1}{x}\right)}$ (viii) $(\log x)^{\sin x} + (x \sin x)^{\frac{1}{x}}$

4. Solve the following :

(i) If $(\sin x)^y = (\sin y)^x$, find $\frac{dy}{dx}$.

(ii) If $(\sin x)^y = (\cos y)^x$, find $\frac{dy}{dx}$.

(iii) If $y = x^y$ show that $\frac{dy}{dx} = \frac{y^2}{x(1-y \log x)}$.

(iv) If $x^y + y^x = \log a$, find $\frac{dy}{dx}$.

5. If $y = \sin^{-1} x$ then show that $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 0$.

6. If $y = (\sin^{-1} x)^2$ then prove that $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 2$.

7. If $y = (\tan^{-1} x)^2$ then show that $(1+x^2)^2 y_2 + 2x(1+x^2)y_1 - 2 = 0$.

8. If $y = \log(x + \sqrt{x^2 + 1})$ then show that $(x^2 + 1) \frac{d^2y}{dx^2} + x \frac{dy}{dx} = 0$.

9. If $y = e^{m \sin^{-1} x}$ then show that $(1-x^2)y_2 - xy_1 - m^2y = 0$.

10. If $y = e^{m \tan^{-1} x}$ prove that $(1+x^2)^2 y_2 + 2x(1+x^2)y_1 - m^2y = 0$.

11. If $u = x^y$, $v = y^x$ and quantity y remains 3 times then quantity x . Find that amongst quantities u and v which changes more rapidly with respect to quantity x when $x = 1$. (Take $\log_e 3 = 1.09$).

12. If $y = (x + \sqrt{x^2 + 1})^m$ then prove that $(x^2 + 1) \frac{d^2y}{dx^2} + x \frac{dy}{dx} - m^2y = 0$.

13. If $y = \cos(2 \cos^{-1} x)$ then prove that $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + 4y = 0$.

14. If $y = \sin(m \sin^{-1} x)$ then show that $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + m^2y = 0$

15. Given $f(x) = (\sin x)^x$, $g(x) = x^{\sin x}$. Find that amongst $f(x)$ and $g(x)$, which function changes less rapidly with respect to x when $x = \frac{\pi}{4}$. Also find the difference between $f'(x)$ and $g'(x)$.

(Take $\frac{1}{\sqrt{2}} = 0.7$, $\frac{\pi}{4} = 0.8$, $\log_e\left(\frac{1}{\sqrt{2}}\right) = -0.4$, $\log_e\left(\frac{\pi}{4}\right) = -0.2$, $(0.7)^{0.8} = 0.8$, $(0.8)^{0.7} = 0.9$)

16. Differentiate the following w.r.t. as indicated :

(i) $\cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$ w.r.t. $\tan^{-1} x$

(ii) $\tan^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right)$ w.r.t. $\sin^{-1} x$

(iii) $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$ w.r.t. $\tan^{-1} x$

(iv) $\cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$ w.r.t. $\tan^{-1} x$

(v) $\tan^{-1}\left(\frac{2x}{1-x^2}\right)$ w.r.t. $\tan^{-1} x$

(vi) $\tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$ w.r.t. $\tan^{-1}\left(\frac{2x}{1-x^2}\right)$

(vii) $\tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$ w.r.t. $\tan^{-1} x$

(viii) $\tan^{-1}\left(\frac{\sqrt{1+a^2x^2}-1}{ax}\right)$ w.r.t. $\tan^{-1} ax$

(ix) $\tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$ w.r.t. $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$

(x) $\cos^2 x$ w.r.t. $e^{\sin x}$

Applications of Derivatives

Multiple Choice Questions (1 Marks)

1 Rate of change of perimeter of a square with respect to its side is :
(a)2 (b)1 (c)4 (d)3

2 Radius of a circle is increasing at the rate of 2 m/s . Rate of change of its circumference is :
(a) $4\pi \text{ m/s}$ (b) 2 m/s (c) $2\pi \text{ m/s}$ (d) 4 m/s

3 Radius of a sphere is increasing at the rate of 5 m/s . Rate of change of its surface area, when radius is 4 m , is
(a) $120\pi \text{ m}^2/\text{s}$ (b) $160\pi \text{ m}^2/\text{s}$ (c) $32\pi \text{ m}^2/\text{s}$ (d) $80\pi \text{ m}^2/\text{s}$

4 $f(x) = \sin x$ is strictly decreasing in the interval :
(a) $(\frac{\pi}{2}, \pi)$ (b) $(\pi, \frac{3\pi}{2})$ (c) $(0, \frac{\pi}{2})$ (d) $(\frac{\pi}{2}, \frac{3\pi}{2})$

5 $f(x) = \cos x$ is strictly increasing in the interval :
(a) $(\frac{\pi}{2}, \pi)$ (b) $(\pi, \frac{3\pi}{2})$ (c) $(0, \frac{\pi}{2})$ (d) $(\frac{\pi}{2}, \frac{3\pi}{2})$

6 $f(x) = x^2$ strictly increases on :
(a) $(0, \infty)$ (b) $(-\infty, 0)$ (c) $(-7, -3)$ (d) $(-\infty, -3)$

7 If f is differentiable at critical points then the value of derivative of f at critical point is :
(a)1 (b)-1 (c)0 (d)2

8 On the curve $y = f(x)$ if $f'(a) = 0$ then $x = a$ is called a
(a)Practical point on the curve (b)Critical point on the curve
(c)Maximum point on the curve (d)Minimum point on the curve

9 On the curve $y = f(x)$ if $f'(a) = 0$ and $f''(a) < 0$ then $x = a$ is point of
(a)Maxima (b)Minima (c)Inflexion (d)infinity

10 On the curve $y = f(x)$ if $f'(a) = 0$ and $f''(a) > 0$ then $x = a$ is point of
(a)Maxima (b)Minima (c)Inflexion (d)infinity

2 Marks Questions

1. The volume of spherical balloon is increasing at the rate of 25 c.c./s . Find the rate of change of its surface area at the instant when its radius is 5 cm.
2. The side of square sheet is increasing at the rate of 3 cm/s. At what rate is the area increasing when the side is 10 cm long?
3. The side of square sheet is increasing at the rate of 5 cm/s. At what rate is the perimeter increasing when the side is 7 cm long?
4. The radius of spherical soap bubble is increasing at the rate of 0.2 cm/s. Find the rate of change of its volume when its radius is 4 cm.
5. The radius of spherical soap bubble is increasing at the rate of 0.8 cm/s. Find the rate of change of its surface area when the radius is 5 cm.
6. The edge of a cube is decreasing at the rate of 2 cm/s . Find the rate of change of its volume when the length of edge of the cube is 5 cm.
7. The edge of a cube is decreasing at the rate of 2 cm/s . Find the rate of change of its surface area when the length of edge is 6 cm.
8. Find the critical points of the following functions :

(a) $f(x) = x^3 + 2x^2 - 1$ (b) $f(x) = 30 - 24x + 15x^2 - 2x^3$ (c) $f(x) = 20 - 12x + 9x^2 - 2x^3$
 (d) $f(x) = 17 - 18x + 12x^2 - 2x^3$ (e) $f(x) = 20 - 9x + 6x^2 - x^3$ (f) $f(x) = 6 + 12x + 3x^2 - 2x^3$
 (g) $f(x) = 2x^3 - 15x^2 + 36x + 1$ (h) $f(x) = x^3 - 6x^2 + 9x + 8$ (i) $f(x) = 2x^3 - 12x^2 + 18x + 5$

9. Determine the intervals in which the following functions are increasing or decreasing :

(a) $f(x) = x^3 + 2x^2 - 1$ (b) $f(x) = 30 - 24x + 15x^2 - 2x^3$ (c) $f(x) = 20 - 12x + 9x^2 - 2x^3$
 (d) $f(x) = 17 - 18x + 12x^2 - 2x^3$ (e) $f(x) = 20 - 9x + 6x^2 - x^3$ (f) $f(x) = 6 + 12x + 3x^2 - 2x^3$
 (g) $f(x) = 2x^3 - 15x^2 + 36x + 1$ (h) $f(x) = x^3 - 6x^2 + 9x + 8$ (i) $f(x) = 2x^3 - 12x^2 + 18x + 5$

6 Marks Questions

- Find the volume of the biggest right circular cone which is inscribed in a sphere of radius 9 cm.
- Prove that the height of a right circular cylinder of maximum volume, which is inscribed in a sphere of radius R , is $\frac{2R}{\sqrt{3}}$.
- Show that of all the rectangles inscribed in a given fixed circle, the square has the maximum area.
- A wire of length 25 m is to be cut into two pieces. One of the pieces is to be made into a square and the other into a circle. What could be the lengths of the two pieces so that the combined area of the square and circle is minimum?
- A wire of length 20 m is to be cut into two pieces. One of the pieces is to be made into a square and the other into an equilateral triangle. What could be the lengths of the two pieces so that the combined area of the square and equilateral triangle is minimum?
- Prove that the perimeter of a right angled triangle of given hypotenuse equal to 5 cm is maximum when the triangle is isosceles.
- Of all rectangles with perimeter 40 cm find the one having maximum area. Also find the area.
- Find the volume of the largest cylinder that can be inscribed in a sphere of radius R .
- Find the volume of largest cone that can be inscribed in a sphere of radius R .
- Show that height of the right circular cylinder of maximum volume that can be inscribed in a sphere of radius 30 cm is $\frac{60}{\sqrt{3}}$ cm.
- A window is in the form of rectangle surmounted by a semi-circle opening. If the perimeter of window is 10 cm, find the dimensions of the window so as to admit maximum possible light through the whole opening.
- Show that the height of a closed cylinder of given volume and least surface area is equal to its diameter.

INTEGRALS

Fill in the Blanks(1 Marks)

1) $\int_0^3 dx = \underline{\hspace{2cm}}$

2) $\int \sec^2 x dx = \underline{\hspace{2cm}}$

3) $\int_0^3 3x^2 dx = \underline{\hspace{2cm}}$

4) $\int_0^5 2x dx = \underline{\hspace{2cm}}$

5) Integration is process of differentiation.

6) $\int_{-a}^a f(x) dx = 0$ if f is function.

7) $\int \frac{dx}{1+x^2} = \underline{\hspace{2cm}}$

8) $\int_0^{\pi/2} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx = \underline{\hspace{2cm}}$

9) $\int \frac{1}{x+3} dx = \underline{\hspace{2cm}}$

10) $\int \sec x \tan x dx = \underline{\hspace{2cm}}$

Multiple Choice Questions(1 Marks)

1 $\int_0^{\pi/2} \frac{\sin^{1/2} x}{\sin^{1/2} x + \cos^{1/2} x} dx$ is equal to :

(a) 0

(b) $\frac{\pi}{2}$

(c) $\frac{\pi}{3}$

(d) $\frac{\pi}{4}$

2 $\int_0^{\pi/2} \frac{\sin^{3/2} x}{\sin^{3/2} x + \cos^{3/2} x} dx$ is equal to :

(a) 0

(b) $\frac{\pi}{2}$

(c) $\frac{\pi}{3}$

(d) $\frac{\pi}{4}$

3 $\int \frac{dx}{2x+3}$ equals:

(a) $\log|2x+3| + c$

(b) $\log|2x-3| + c$

(c) $\frac{\log|2x+3|}{3} + c$

(d) $\frac{\log|2x+3|}{2} + c$

4 $\int \frac{dx}{2x-5}$ equals:

(a) $\log|2x-5| + c$

(b) $\log|2x+5| + c$

(c) $\frac{\log|2x-5|}{5} + c$

(d) $\frac{\log|2x-5|}{2} + c$

5 $\int_{-1}^1 x^3 \cos x dx$ equals:

(a) 0

(b) 1/4

(c) π

(d) none of these

6 $\int_0^{\pi/2} \frac{\sin^{1/2} x}{\sin^{1/2} x + \cos^{1/2} x} dx$ is equal to :

(a) 0

(b) $\frac{\pi}{2}$

(c) $\frac{\pi}{3}$

(d) $\frac{\pi}{4}$

7 $\int_{-2}^2 x^3 dx$ is equal to :

(a) 0

(b) 4

(c) $16/3$

(d) $\frac{\pi}{4}$

8 $\int_{-1}^1 x \sin^2 x dx$ is equal to

(a) 0

(b) $\frac{1}{2}$

(c) $\frac{1}{3}$

(d) -1

9 $\int_0^1 \frac{dx}{1+x^2}$ is
 (a) $\frac{\pi}{2}$ (b) $\frac{\pi}{3}$ (c) $\frac{\pi}{4}$ (d) $\frac{\pi}{6}$

10 $\int_{\pi/6}^{\pi/3} \frac{\cos^4 x}{\sin^4 x + \cos^4 x} dx$ is equal to
 (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{6}$ (c) $\frac{\pi}{12}$ (d) $\frac{\pi}{2}$

2 Marks Questions

1. Evaluate the following :

$$\begin{array}{llll}
 \text{(a)} \int \frac{(x-4)^3}{x^2} dx & \text{(b)} \int \frac{dx}{1-\sin x} & \text{(c)} \int \frac{dx}{1+\cos x} & \text{(d)} \int \frac{dx}{1+\sin x} \\
 \text{(e)} \int \frac{dx}{1-\cos x} & \text{(f)} \int \frac{e^x-1}{e^x+1} dx & \text{(g)} \int \frac{(\tan^{-1} x)^2}{1+x^2} dx & \text{(h)} \int \frac{\sec^2(2 \tan^{-1} x)}{1+x^2} dx \\
 \text{(i)} \int \frac{\sin(\tan^{-1} x)}{1+x^2} dx & \text{(j)} \int \frac{dx}{x^2+8x-9} & \text{(k)} \int \frac{dx}{\sqrt{x^2-5x+7}} & \text{(l)} \int \frac{dx}{\sqrt{x^2+4x+7}} \\
 \text{(m)} \int \frac{dx}{x^2+6x+5} & \text{(n)} \int \frac{dx}{x^2-6x+18} & \text{(o)} \int x\sqrt{x+2} dx & \text{(p)} \int \frac{3-2 \sin x}{\cos^2 x} dx
 \end{array}$$

2. Compute the following :

$$\begin{array}{llll}
 \text{(a)} \int \frac{\log x}{x} dx & \text{(b)} \int \frac{e^{\tan^{-1} x}}{1+x^2} dx & \text{(c)} \int \frac{2x}{1+x^2} dx & \text{(d)} \int \frac{x^2}{1+x^3} dx \\
 \text{(e)} \int \frac{6x-8}{3x^2-8x+5} dx & \text{(f)} \int \frac{2x+9}{x^2+9x+20} dx & \text{(g)} \int e^x \left(\tan^{-1} x + \frac{1}{1+x^2} \right) dx
 \end{array}$$

4 Marks Questions

3. Integrate the following :

$$\begin{array}{lll}
 \text{(a)} \sin^2 x \cos^3 x & \text{(b)} \cos^2 x \sin^3 x & \text{(c)} \frac{1}{1-\cot x} \\
 \text{(d)} \frac{1}{1+\cot x} & \text{(e)} \frac{1}{1-\tan x} & \text{(f)} \frac{1}{1+\tan x}
 \end{array}$$

4. Evaluate the following integrals :

$$\text{(a)} \int \frac{1-\tan x}{1+\tan x} dx \quad \text{(b)} \int \frac{1+\tan x}{1-\tan x} dx$$

5. Integrate the following functions:

$$\text{(a)} x \sec^2 x \quad \text{(b)} x^2 e^x \quad \text{(c)} x \cos 3x \quad \text{(d)} x \sin x$$

7. Integrate the following functions :

(a) $e^x \sin 2x$ (b) $e^{3x} \cos 5x$ (c) $e^x(\cot x + \log \sin x)$

8. Integrate the following functions :

(a) $\frac{1}{(x+1)(x+2)(x+3)}$ (b) $\frac{1}{x(x-1)(x-2)}$ (c) $\frac{1}{x^3-1}$ (d) $\frac{1}{(1-x)(1+x^2)}$

(e) $\frac{x}{(x-2)(x^2+4)}$ (f) $\frac{1}{x(x^2+2)}$

9. Integrate the following functions :

(a) $\frac{2x}{\sqrt{(x+1)(x-2)}}$ (b) $\frac{4x+5}{\sqrt{x^2+x-3}}$ (c) $\frac{3x+5}{\sqrt{x^2-8x+7}}$

12 Evaluate the following integrals :

(a) $\int_0^{\pi/2} \frac{\sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$ (b) $\int_0^{\pi/2} \frac{1}{1 + \sqrt{\cot x}} dx$

(c) $\int_0^{\pi/2} \frac{\sqrt{\cot x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx$ (e) $\int_0^1 |x - 5| dx$

(f) $\int_{-6}^6 |x + 2| dx$ (h) $\int_0^{\pi/2} \log(\cos x) dx$

13 Prove that (i) $\int \frac{dx}{x^2+a^2} = \log |x + \sqrt{x^2 + a^2}| + C$

(ii) $\int \frac{dx}{\sqrt{a^2-x^2}} = \sin^{-1} \left(\frac{x}{a} \right) + C$

6 Marks Questions

Evaluate the following :

1. $\int \frac{x^2+1}{x^4+1} dx$

2. $\int \frac{x^2}{x^4+1} dx$

3. $\int \frac{1}{x^4+1} dx$

4. $\int_0^{\pi/2} \log \cos x dx$

5. $\int \frac{2x}{(x^2+1)(x^2+4)} dx$

6. $\int \frac{1}{x^3-1} dx$

7. $\int (\sqrt{\cot x} + \sqrt{\tan x}) dx$

8. $\int_0^{\pi} \frac{x dx}{a^2 \cos^2 x + b^2 \sin^2 x}$

APPLICATIONS OF INTEGRALS**2 Marks Questions**

1. Using integration, find the area of the circle :

(i) $x^2 + y^2 = 4$

(ii) $x^2 + y^2 = 9$

(iii) $\frac{x^2}{9} + \frac{y^2}{4} = 1$

(iv) $\frac{x^2}{9} + \frac{y^2}{25} = 1$

(v) $\frac{x^2}{16} + \frac{y^2}{36} = 1$

(vi) $\frac{x^2}{25} + \frac{y^2}{16} = 1$

2. Find the area of the region bounded by $x^2 + y^2 = 16$, $y = x$ in the first quadrant.

3. Find the area of the region bounded by $x^2 + y^2 = 25$, $y = 2x$ in the first quadrant

4. Draw a rough sketch to indicate the region bounded between the curve $y^2 = 4x$, $x = 3$. Also find the area of this region.

5. Find the area of the region bounded by the curves $y^2 = 8x$, $x = 1$, $x = 5$ in the first quadrant.

6. Find the area bounded by the three line $y = x$, $x = 2$ and $x = 5$ in the first quadrant.

7. Find the area bounded by the parabola $y^2 = 16x$ and its latusrectum.

DIFFERENTIAL EQUATIONS

Multiple Choice Questions(1 Marks)

1 Integrating factor of differential equation $\frac{dy}{dx} - \frac{y}{x} = 2x$ is :
 (a) $\frac{1}{x}$ (b) x (c) $\frac{1}{x^2}$ (d) 1

2 Order of differential equation $\frac{d^2y}{dx^2} - \left(\frac{dy}{dx}\right)^3 + y = 0$ is :
 (a) 3 (b) 2 (c) 0 (d) 1

3 Differential equation for the family of the curves $y^2 = kx$ is:
 (a) $\frac{dy}{dx} = 0$ (b) $y + 2x \frac{dy}{dx} = 0$ (c) $y - 2x \frac{dy}{dx} = 0$ (d) $y \frac{dy}{dx} = 1$

4 Integrating factor of differential equation $\frac{dy}{dx} + \frac{y}{x} = 2x$ is:
 (a) $\frac{1}{x}$ (b) x^2 (c) $\frac{1}{x^2}$ (d) x

5 Integrating factor of differential equation $\frac{dy}{dx} + \frac{2y}{x} = 2x$ is:
 (a) $\frac{1}{x}$ (b) x^2 (c) $\frac{1}{x^2}$ (d) x

6 Integrating factor of differential equation $\frac{dy}{dx} + y \sec x = 2x$ is :
 (a) $\sec x + \tan x$ (b) $\sec x \tan x$ (c) $e^{\sec x}$ (d) $e^{\sec x + \tan x}$

7 Integrating factor of differential equation $\frac{dy}{dx} + y = 2x$ is :
 (a) $\frac{1}{x}$ (b) x (c) e^x (d) e^{-x}

8 Order of differential equation $\frac{d^3y}{dx^3} - 4 \left(\frac{d^2y}{dx^2}\right)^4 + y = 0$ is
 (a) 3 (b) 4 (c) 1 (d) 0

9 The number of arbitrary constants in the general solution of a differential equation of second order are
 (a) 1 (b) 2 (c) 3 (d) 4

10 Degree of the differential equation $\frac{d^3y}{dx^3} - \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^4 + y = 0$ is
 (a) 1 (b) 2 (c) 3 (d) 4

Fill Ups(1 Marks)

1) Order of the differential equation $\frac{d^2y}{dx^2} - \left(\frac{dy}{dx}\right)^3 + y = 0$ is _____

2) Degree of the differential equation $\frac{d^2y}{dx^2} - \left(\frac{dy}{dx}\right)^3 + y = 0$ is _____

3) Integrating factor of differential equation $\frac{dy}{dx} + xy = \sin x$ is _____

4) Order and degree (if defined) of a differential equation are always _____ integers.

5) Integrating factor of $\frac{dx}{dy} + Px = Q$ is _____

6) $(x + y)dy - (x - 2y)dx = 0$ is a _____ differential equation.

7) _____ substitution is applied to solve a homogeneous differential equation.

8) There are _____ number of arbitrary constants in the general solution of differential equation of order 3.

9) Differential equation representing the family of curve $y = mx + c$ is given by _____

10) After correct substitution, a homogeneous differential equation becomes _____ type of differential equation.

4 Marks Questions

Solve the following differential equations :

1) $\frac{dy}{dx} = \log x$

2) $\frac{dy}{dx} + \frac{1+y^2}{y} = 0$

3) $\frac{dy}{dx} = \sin^2 y$

4) $\frac{dy}{dx} = e^y \sin x$

5) $\log \frac{dy}{dx} = ax + by$

6) $x^2(y+1)dx + y^2(x-1)dy = 0$

7) $\sec^2 x \tan y dx - \sec^2 y \tan x dy = 0$

8) $xdy + ydx = xydx ; y(1) = 1$

9) $x(xdy - ydx) = ydx ; y(1) = 1$

10) $\frac{dy}{dx} = y \tan x ; y(0) = 1$

11) $\frac{dy}{dx} = y \sin 2x ; y(0) = 1$

12) $(x^2 + xy)dy + (3xy + y^2)dx = 0$

13) $(y^2 - x^2)dy - 3xydx = 0$

14) $2xydx + (x^2 + 2y^2)dy = 0$

15) $x^2dy - (x^2 + xy + y^2)dx = 0$

16) $\cos\left(\frac{dy}{dx}\right) = \frac{1}{9} ; y(0) = 2$

17) $(x^2 + y^2)dx + 2xydy = 0$

18) $(x^2 - 2y^2)dx + xydy = 0$

19) $\frac{dy}{dx} + \frac{y}{x} = e^x$

20) $\frac{dy}{dx} - 4y = e^{2x}$

21) $x \frac{dy}{dx} + y = x^3$

22) $\frac{dy}{dx} + 2y = \sin 5x$

23) $\frac{dy}{dx} + 3y = \cos 2x$

24) $x \frac{dy}{dx} + y = x \log x$

25) $(1 + x^2) \frac{dy}{dx} + y = \tan^{-1} x$

26) $\frac{dy}{dx} = 2x + y ; y(0) = 0$

27) $\frac{dy}{dx} = 4x + y ; y(0) = 1$

28) $x \frac{dy}{dx} + y = x^3 ; y(2) = 1$

29) $xy' - y = \log x ; y(1) = 0$

30) $x \log x \frac{dy}{dx} + y = \frac{2}{x} \log x$

31) $\frac{dy}{dx} = \frac{1-\cos x}{1+\cos x}$

32) $(1 + x^2)dy + 2xydx = \cot x dx$

33) $x \frac{dy}{dx} + 2y = x^2 \log x$

34) $x^2dy - (3x^2 + xy + y^2)dx = 0 ; y(1) = 1.$

Vector Algebra

Fill in the Blanks(1 Mark)

- 1) Scalar product of two perpendicular vectors is always equal to _____.
- 2) Vector product of two collinear vectors is always equal _____.
- 3) $|\vec{a} \cdot \vec{b}| \leq |\vec{a}| |\vec{b}|$ is called _____ inequality.
- 4) $|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$ is called _____ inequality.
- 5) $\hat{i} \times \hat{i} = \hat{j} \times \hat{j} = \hat{k} \times \hat{k} = \text{_____}$.
- 6) $\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = \text{_____}$
- 7) $\vec{a} \times \vec{b}$ is _____ to both the vectors \vec{a} and \vec{b} .
- 8) Area of a parallelogram can be calculated by using _____ product of two vectors.

Multiple Choice Questions(1 Mark)

- 1) If $\vec{a} \cdot \vec{b} = |\vec{a} \times \vec{b}|$ then angle between vector \vec{a} and vector \vec{b} is :

(a) $\frac{\pi}{2}$	(b) $\frac{\pi}{6}$	(c) $\frac{\pi}{4}$	(d) $\frac{\pi}{3}$
---------------------	---------------------	---------------------	---------------------
- 2) Magnitude of the vector $\frac{1}{\sqrt{3}} \hat{i} + \frac{1}{\sqrt{3}} \hat{j} + \frac{1}{\sqrt{3}} \hat{k}$ is :

(a) -1	(b) 1	(c) 0	(d) $\frac{1}{3}$
--------	-------	-------	-------------------
- 3) If $\sqrt{3} \vec{a} \cdot \vec{b} = |\vec{a} \times \vec{b}|$ then angle between vector \vec{a} and vector \vec{b} is :

(a) $\frac{\pi}{2}$	(b) $\frac{\pi}{6}$	(c) $\frac{\pi}{4}$	(d) $\frac{\pi}{3}$
---------------------	---------------------	---------------------	---------------------
- 4) If $\vec{a} \cdot \vec{b} = \sqrt{3} |\vec{a} \times \vec{b}|$ then angle between vector \vec{a} and vector \vec{b} is :

(a) $\frac{\pi}{2}$	(b) $\frac{\pi}{6}$	(c) $\frac{\pi}{4}$	(d) $\frac{\pi}{3}$
---------------------	---------------------	---------------------	---------------------
- 5) If $\vec{a} \cdot \vec{b} = 0$ then angle between vector \vec{a} and vector \vec{b} is :

(a) $\frac{\pi}{2}$	(b) $\frac{\pi}{6}$	(c) $\frac{\pi}{4}$	(d) $\frac{\pi}{3}$
---------------------	---------------------	---------------------	---------------------
- 6) Name of the inequality $|\vec{a} \cdot \vec{b}| \leq |\vec{a}| |\vec{b}|$ is :

(a) Cauchy-Schwartz Inequality	(b) Triangle Inequality
(c) Rolle's Inequality	(d) Lagrange's Inequality
- 7) Magnitude of vector $\vec{a} = 3 \hat{i} + \hat{j} + \hat{k}$ is :

(a) 3	(b) $\sqrt{10}$	(c) $\sqrt{11}$	(d) $\sqrt{12}$
-------	-----------------	-----------------	-----------------

8 Projection of $\vec{a} = 3\hat{i} + \hat{j} + \hat{k}$ on $\vec{b} = \hat{i} - 2\hat{j} - \hat{k}$ is :

(a) $\frac{2}{\sqrt{6}}$

(b) 0

(c) $\frac{1}{\sqrt{6}}$

(d) $\sqrt{6}$

9 If \vec{a} is a non-zero vector then $|\vec{a} \times \vec{a}|$ is equal to

(a) $|\vec{a}|$

(b) $|\vec{a}|^2$

(c) 1

(d) 0

10 If $\vec{a} = \hat{i} + 2\hat{j} - 3\hat{k}$ and $\vec{b} = 2\hat{i} - 2\hat{j} - \hat{k}$ then $\vec{a} \cdot \vec{b}$ is equal to

(a) 1

(b) 0

(c) -1

(d) 3

2 Marks Questions

1. Adjacent sides of a parallelogram are given by $\hat{i} + 2\hat{j} - \hat{k}$ and $3\hat{i} - \hat{j} + 5\hat{k}$. Find a unit vector along a diagonal of the parallelogram.
2. Adjacent sides of a parallelogram are given by $6\hat{i} - \hat{j} + 5\hat{k}$ and $\hat{i} + 5\hat{j} - 2\hat{k}$. Find the area of parallelogram.
3. Find the area of triangle whose sides are given by the vectors $\hat{i} - 2\hat{j} + \hat{k}$ and $4\hat{i} + \hat{j} - 7\hat{k}$.
4. Find the value of p if the vectors $p\hat{i} + \hat{j} + 4\hat{k}$ and $2\hat{i} - \hat{j} + 3\hat{k}$ are perpendicular to each other.
5. Find a vector of magnitude 8 units along $\vec{a} = 2\hat{i} - 4\hat{j} + \hat{k}$
6. Find a unit vector along $\vec{a} = 5\hat{i} + 3\hat{j} - 4\hat{k}$
7. If $\vec{a} = 2\hat{i} - 4\hat{j} + \hat{k}$, $\vec{b} = 3\hat{i} - \hat{j} - 5\hat{k}$ then find $|\vec{a} \times \vec{b}|$.
8. Find the projection of $\vec{a} = 2\hat{i} - 4\hat{j} + \hat{k}$ on $\vec{b} = 3\hat{i} - \hat{j} - 5\hat{k}$.
9. Find the area of parallelogram whose diagonals are given by vectors:

(i) $\vec{a} = 2\hat{i} + \hat{j} + \hat{k}$ & $\vec{b} = \hat{i} - \hat{k}$

(ii) $\vec{a} = \hat{i} + \hat{j} - 4\hat{k}$ & $\vec{b} = \hat{i} + 8\hat{j} + 2\hat{k}$

10. Find the angle between the following vectors:

(i) $\vec{a} = \hat{i} - 2\hat{j} + 4\hat{k}$ and $\vec{b} = -2\hat{i} + 4\hat{j} - 2\hat{k}$

(ii) $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$ and $\vec{b} = 3\hat{i} + 2\hat{j} - 7\hat{k}$

3/4 Marks Questions

- For any two vectors \vec{a} and \vec{b} prove that $|\vec{a} \cdot \vec{b}| \leq |\vec{a}| |\vec{b}|$. Also write the name of inequality.
- For any two vectors \vec{a} and \vec{b} prove that $|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$. Also write the name of inequality.
- Find the area of triangle whose vertices are :
 - $A(2, 3, 5), B(3, 5, 8), C(2, 7, 8)$
 - $A(1, 2, 4), B(3, 1, -2), C(4, 3, 1)$
 - $P(1, 1, 1), Q(1, 2, 3), R(2, 3, 1)$
- Let $\vec{a}, \vec{b}, \vec{c}$ be three vectors such that $|\vec{a}| = 3, |\vec{b}| = 4, |\vec{c}| = 5$ and each one of them is perpendicular to the sum of the other two, find $|\vec{a} + \vec{b} + \vec{c}|$.
- If $\vec{a} = 5\hat{i} + \hat{j} - 2\hat{k}, \vec{b} = 7\hat{i} + 2\hat{j} - 3\hat{k}, \vec{c} = 2\hat{i} - 9\hat{j} - \hat{k}$ find a vector of magnitude 7 units parallel to the vector $2\vec{a} - \vec{b} + 3\vec{c}$.
- If $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}, \vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}, \vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$, then find a vector \vec{d} which is perpendicular to both \vec{a} and \vec{b} and $\vec{c} \cdot \vec{d} = 15$.

Three Dimensional Geometry

Multiple Choice Questions(1 Marks)

- 1 Direction ratios of straight line $\vec{r} = \hat{i} - 4\hat{j} + 5\hat{k} + s(2\hat{i} - 3\hat{j} + 2\hat{k})$ are :
 (a) $< 2, 3, 2 >$ (b) $< 2, -3, -2 >$ (c) $< -2, -3, 2 >$ (d) $< 2, -3, 2 >$
- 2 Direction ratios of line given by $\frac{x-1}{3} = \frac{2y+6}{12} = \frac{1-z}{-7}$ are :
 (a) $< 3, 12, -7 >$ (b) $< 3, -6, 7 >$ (c) $< 3, 6, 7 >$ (d) $< 3, 6, -7 >$
- 3 Direction ratios of a line passing through the points $(-2, 1, 0)$ & $(3, 2, 1)$ are
 (a) $< 5, 1, 1 >$ (b) $< -5, 1, -1 >$ (c) $< 5, -1, 1 >$ (d) $< -5, -1, 1 >$
- 4 Which of the following sets of points are collinear :
 (a) $(1, 3, -4), (1, -2, 7)$ & $(3, 8, -11)$ (b) $(2, 3, -4), (2, -2, 3)$ & $(3, 5, -11)$
 (c) $(2, 3, -4), (1, -2, 3)$ & $(3, 8, -11)$ (d) $(2, 3, -4), (1, -2, 3)$ & $(2, 8, 11)$
- 5 Vector equation of the line $\frac{x+4}{5} = \frac{y-5}{3} = \frac{z-8}{-3}$ is
 (a) $\vec{r} = 4\hat{i} - 5\hat{j} - 8\hat{k} + \mu(5\hat{i} + 3\hat{j} - 3\hat{k})$ (b) $\vec{r} = -4\hat{i} + 5\hat{j} + 8\hat{k} + \mu(5\hat{i} + 3\hat{j} - 3\hat{k})$
 (c) $\vec{r} = 5\hat{i} + 3\hat{j} - 3\hat{k} + \mu(4\hat{i} - 5\hat{j} - 8\hat{k})$ (d) $\vec{r} = 5\hat{i} + 3\hat{j} - 3\hat{k} + \mu(-4\hat{i} + 5\hat{j} - 8\hat{k})$
- 6 Cartesian equation of the line $\vec{r} = 7\hat{i} - 5\hat{j} + 3\hat{k} + \mu(9\hat{i} - \hat{j} + 6\hat{k})$ is
 (a) $\frac{x+9}{7} = \frac{y-1}{-5} = \frac{z+6}{3}$ (b) $\frac{x-9}{7} = \frac{y+1}{-5} = \frac{z+6}{3}$ (c) $\frac{x+7}{9} = \frac{y-5}{-1} = \frac{z+3}{6}$ (d) $\frac{x-7}{9} = \frac{y+5}{-1} = \frac{z-3}{6}$
- 7 Angle between the lines $\frac{x+1}{2} = \frac{y-5}{-1} = \frac{z}{1}$ and $\frac{x}{3} = \frac{y+7}{5} = \frac{z-8}{-1}$ is
 (a) $\pi/3$ (b) $\pi/2$ (c) $\pi/6$ (d) 0
- 8 Direction cosines of a line making equal angles with coordinate axes are
 (a) $< \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} >$ (b) $< \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} >$ (c) $< 1, 1, 1 >$ (d) $< 0, 0, 0 >$
- 9 Direction ratios of a line making equal angles with coordinate axes are
 (a) $< \frac{1}{\sqrt{3}}, \frac{-1}{\sqrt{3}}, \frac{1}{\sqrt{3}} >$ (b) $< \frac{-1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} >$ (c) $< 1, 1, 1 >$ (d) $< 0, 0, 0 >$
- 10 If direction ratios of a line are $< 9, -6, 2 >$ then its direction cosines are
 (a) $< \frac{9}{\sqrt{117}}, \frac{-6}{\sqrt{117}}, \frac{2}{\sqrt{117}} >$ (b) $< \frac{9}{\sqrt{121}}, \frac{-6}{\sqrt{121}}, \frac{2}{\sqrt{121}} >$ (c) $< \frac{9}{7}, \frac{-6}{7}, \frac{2}{7} >$ (d) $< \frac{9}{5}, \frac{-6}{5}, \frac{2}{5} >$

Fill in the Blanks(1 Mark)

- 1) Direction cosine of y – axis are _____.
- 2) The equation of a line parallel to the vector $2\hat{i} - \hat{j} + 3\hat{k}$ and passing through the point $(5, -1, 3)$ is _____.
- 3) The angle between two lines with direction ratios $< l, m, n >$ and $< a, b, c >$ is given by _____.
- 4) Direction ratios of a line which passes through the points $(1, 2, 3)$ and $(4, 1, -2)$ are _____.
- 5) Direction ratios of a line which makes equal angles with coordinate axes are _____.
- 6) Direction ratios of line are $< 1, 0, 0 >$, then its direction cosines are _____.
- 7) Vector equation of a line which passes through the origin and makes equal angles with the coordinate axes is _____.
- 8) Cartesian equation of the line $\vec{r} = 3\hat{j} - \hat{i} + 2\hat{k} + t(5\hat{k} + \hat{i} - 7\hat{j})$ is _____.

2/3 Marks Questions

- Find the equation of a line which passes through the points $(3, 6, -7)$ and $(5, -1, 4)$.
- Find the direction cosines of a line passing through the points $(7, -1, 2)$ and $(3, 4, -7)$.
- Find the direction ratios and direction cosines of a line which makes equal angles with the coordinate axes.
- Find the direction cosines of sides of a triangle whose vertices are $(1, 2, -3)$, $(9, -3, 7)$ and $(5, 3, -2)$.
- Find the angle between the lines :

(i) $\vec{r} = 3\hat{i} + 8\hat{j} + 3\hat{k} + \mu(3\hat{i} - \hat{j} + \hat{k})$ & $\vec{r} = -3\hat{i} - 7\hat{j} + 6\hat{k} + \lambda(-3\hat{i} + 2\hat{j} + 4\hat{k})$

(ii) $\vec{r} = 2\hat{i} - \hat{j} - \hat{k} + \mu(3\hat{i} - 5\hat{j} + 2\hat{k})$ & $\vec{r} = \hat{i} + 2\hat{j} + \hat{k} + \lambda(\hat{i} - \hat{j} + \hat{k})$

(iii) $\frac{x-1}{1} = \frac{y-2}{-1} = \frac{z-1}{1}$ & $\frac{x-2}{2} = \frac{y+1}{1} = \frac{z+1}{2}$

(iv) $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ & $\frac{x-2}{3} = \frac{y-4}{5} = \frac{z-5}{5}$

- Find the value of m if the lines $\frac{x+2}{3} = \frac{y-1}{2m} = \frac{z-2}{7}$ and $\frac{x-3}{4} = \frac{y-2}{7} = \frac{z+5}{8m}$ are perpendicular to each other.

6/4 Marks Questions

- Find the shortest distance between the following pairs of lines :

(i) $\vec{r} = 3\hat{i} + 8\hat{j} + 3\hat{k} + \mu(3\hat{i} - \hat{j} + \hat{k})$ & $\vec{r} = -3\hat{i} - 7\hat{j} + 6\hat{k} + \lambda(-3\hat{i} + 2\hat{j} + 4\hat{k})$

(ii) $\vec{r} = 2\hat{i} - \hat{j} - \hat{k} + \mu(3\hat{i} - 5\hat{j} + 2\hat{k})$ & $\vec{r} = \hat{i} + 2\hat{j} + \hat{k} + \lambda(\hat{i} - \hat{j} + \hat{k})$

(iii) $\frac{x-1}{1} = \frac{y-2}{-1} = \frac{z-1}{1}$ & $\frac{x-2}{2} = \frac{y+1}{1} = \frac{z+1}{2}$

(iv) $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ & $\frac{x-2}{3} = \frac{y-4}{5} = \frac{z-5}{5}$

(v) $\frac{3-x}{2} = \frac{8-2y}{-10} = \frac{z-1}{1}$ & $\frac{3x-6}{9} = \frac{5-y}{1} = \frac{4-2z}{8}$

LINEAR PROGRAMMING

Multiple Choice Questions(1 Mark)

- 1 All points of feasible region are :
(a)infeasible solutions (b)feasible solutions (c)optimal solutions (d)none of these
- 2 Corner points of the feasible region are
(a)optimal solutions (b)useless points (c)infeasible solutions (d)none of these
- 3 Common area for each constraint is called :
(a)infeasible region (b)feasible region (c)useless area (d)none of these
- 4 The objective function of linear programming problem is :
(a)a linear function to be minimized or maximized
(b)a linear expression representing constraints
(c)a quadratic equation representing the corner points of feasible region
(d)represents the area of the feasible region
- 5 Minimum value of $Z = 4x + 3y$ subject to the constraints $x + y \leq 4$, $x, y \geq 0$ is
(a)16 (b)12 (c)10 (d)20
- 6 The corner points of bounded feasible region of linear programming problem are $(0, 0)$, $(0, 4)$, $(8, 0)$ and $(\frac{20}{3}, \frac{4}{3})$ then one of the constraints is :
(a) $2x + 5y \leq 20$ (b) $2x + 5y \geq 20$ (c) $5x + 2y \leq 20$ (d) $5x + 2y \geq 20$
- 7 Minimum value of $Z = 5x + 3y + 2$ subject to the constraints $x + y \leq 7$, $x, y \geq 0$ is
(a)37 (b)35 (c)21 (d)23
- 8 Constraints of LPP are :
(a)Always quadratic (b)Always linear
(c)May be linear or quadratic depending on the problem (d)May be cubic some times
- 9 Objective function of LPP is
(a)Always quadratic (b)Always linear
(c)May be linear or quadratic depending on the problem (d)May be cubic some times
- 10 Minimum value of $Z = 5x + 3y + 2$ subject to the constraints $x + y \leq 7$, $x, y \geq 0$ on the point
(a) $(7, 0)$ (b) $(0, 7)$ (c) $(3, 4)$ (d) $(4, 3)$

4 Marks Questions

Solve the following LPP graphically:

1. Maximize & Minimize :

- (i) $Z = 10x + 7y$ subject to the constraints $3x + y \leq 9$, $3x + 2y \leq 12$, $x, y \geq 0$.
- (ii) $Z = x + 2y$ subject to the constraints $7x + 3y \leq 21$, $x + y \geq 3$, $x - y \leq 0$, $x, y \geq 0$.
- (iii) $Z = 4x + 2y$ subject to the constraints $8x + 9y \leq 72$, $4x + y \geq 8$, $2x - y \geq 0$, $x, y \geq 0$.
- (iv) $Z = 5x + 7y$ subject to the constraints $x + y \geq 4$, $x + 3y \leq 12$, $x - 2y \geq 0$, $x, y \geq 0$.
- (v) $Z = 3x + 6y$ subject to the constraints $x + y \leq 6$, $2x + y \geq 6$, $2x - y \leq 0$, $x, y \geq 0$.
- (vi) $Z = 4x + 5y$ subject to the constraints $x + y \leq 6$, $2x + y \geq 6$, $x - y \geq 0$, $x, y \geq 0$.
- (vii) $Z = 8x + y$ subject to the constraints $x + y \leq 8$, $2x + y \geq 8$, $x - 2y \leq 0$, $x, y \geq 0$.
- (viii) $Z = 3x + 7y$ subject to the constraints $x + y \leq 10$, $x + 2y \geq 6$, $3x - y \leq 9$, $x, y \geq 0$.
- (ix) $Z = 8x + 5y - 2$ subject to the constraints $x + y \leq 10$, $x + 2y \geq 6$, $3x - y \geq 9$, $x, y \geq 0$
- (x) $Z = 7x + 5y - 1$ subject to the constraints $x + y \leq 10$, $x + y \geq 5$, $x - y \leq 0$, $x, y \geq 0$

PROBABILITY

Multiple Choice Questions(1 Mark)

1 If $P(A) = \frac{1}{2}$, $P(B) = \frac{3}{8}$ and $P(A \cap B) = \frac{1}{5}$ then $P(A|B)$ is equal to :
 (a) $\frac{2}{5}$ (b) $\frac{8}{15}$ (c) $\frac{2}{3}$ (d) $\frac{5}{8}$

2 If A and B are independent events and $P(A) = \frac{1}{2}$, $P(B) = \frac{3}{8}$ then $P(A \cap B)$ is equal to :
 (a) $\frac{3}{4}$ (b) $\frac{3}{8}$ (c) $\frac{3}{16}$ (d) $\frac{1}{16}$

3 If $P(A) = \frac{1}{2}$, $P(B) = \frac{3}{8}$ and $P(A \cap B) = \frac{1}{5}$ then $P(B|A)$ is equal to :
 (a) $\frac{2}{5}$ (b) $\frac{8}{15}$ (c) $\frac{2}{3}$ (d) $\frac{5}{8}$

4 Probability of getting even prime number on both dice ,on a throw of a pair of die, is :
 (a) $1/6$ (b) $2/35$ (c) $1/36$ (d) $5/36$

5 If $P(A) = \frac{1}{2}$, $P(B) = \frac{3}{8}$ and $P(A \cup B) = \frac{4}{5}$ then $P(A|B)$ is equal to :
 (a) $\frac{1}{5}$ (b) $\frac{8}{15}$ (c) $\frac{2}{3}$ (d) $\frac{5}{8}$

6 If E is any event then $P(E)$ belongs to the interval :
 (a) $(1, 10)$ (b) $(0, 1)$ (c) $[0, 1]$ (d) $[10, 20]$

7 If $P(E) = \frac{5}{7}$ then $P(\text{not } E)$ is
 (a) $\frac{5}{7}$ (b) $\frac{7}{5}$ (c) $\frac{7}{2}$ (d) $\frac{2}{7}$

8 A coin is marked with head on both sides then on tossing the coin probability of getting head is :
 (a) $\frac{1}{2}$ (b) 0 (c) 1 (d) 2

9 Probability of getting an ace card on drawing one card from a well shuffled deck of 52 cards is :
 (a) $1/13$ (b) $1/4$ (c) $1/52$ (d) none of these

10 Probability of 53 Mondays in a leap year is
 (a) $\frac{2}{53}$ (b) $\frac{2}{7}$ (c) $\frac{1}{53}$ (d) $\frac{1}{7}$

11 Probability of 53 Mondays in a non-leap year is
 (a) $\frac{2}{53}$ (b) $\frac{2}{7}$ (c) $\frac{1}{53}$ (d) $\frac{1}{7}$

12 If E is any event then $P(\text{not } E)$ belongs to the interval :
 (a) $(1, 10)$ (b) $(0, 1)$ (c) $[0, 1]$ (d) $[10, 20]$

13 If three coins are tossed once, then getting at least one heads is
 (a) $\frac{3}{8}$ (b) $\frac{7}{8}$ (c) $\frac{5}{8}$ (d) $\frac{1}{2}$

14 There are 3 red balls, 4 white balls and 7 blue balls in a bag. One ball is drawn at random from the bag.
 Probability of drawing a white ball is
 (a) $\frac{2}{7}$ (b) $\frac{3}{14}$ (c) $\frac{7}{14}$ (d) 0

15 There are 3 red balls, 4 white balls and 7 blue balls in a bag. One ball is drawn at random from the bag.
 Probability of drawing a green ball is
 (a) $\frac{2}{7}$ (b) $\frac{3}{14}$ (c) $\frac{7}{14}$ (d) 0

16 If E and F are independent events , then
 (a) $P(E \cup F) = P(E) + P(F)$ (c) $P(E \cap F) = P(E) + P(F)$
 (c) $P(E \cap F) = P(E) \cdot P(F)$ (d) $P(E \cap F) = 0$

Fill in the Blanks(1 Mark)

- 1) If $P(A) = \frac{1}{5}$ then $P(\text{not } A) = \underline{\hspace{2cm}}$
- 2) In a throw of a pair of dice probability of getting a doublet is $\underline{\hspace{2cm}}$
- 3) Probability of occurrence of sure event = $\underline{\hspace{2cm}}$
- 4) Probability of occurrence of impossible event = $\underline{\hspace{2cm}}$
- 5) $P(A \cup B) = P(A) + P(B) - \underline{\hspace{2cm}}$
- 6) $P(A) + \underline{\hspace{2cm}} = 1$
- 7) If A and B are independent events then $P(A \cap B) = \underline{\hspace{2cm}}$
- 8) If $P(A) = \frac{1}{2}$ and $P(B) = 0$ then $P(A|B)$ is $\underline{\hspace{2cm}}$
- 9) If a dice is tossed once then probability of getting an odd prime number is $\underline{\hspace{2cm}}$
- 10) Probability of any event is (numerically) always less than or equal to $\underline{\hspace{2cm}}$

4 Marks Questions

1. If $P(A) = \frac{6}{11}$, $P(B) = \frac{5}{11}$ and $P(A \cup B) = \frac{7}{11}$ then find : $P(A \cap B)$, $P(A|B)$ & $P(B|A)$
2. If $P(E) = 0.45$, $P(F) = 0.55$ & $P(E \cup F) = 0.75$ then find $P(E \cap F)$ & $P(E|F)$.
3. If A & B are independent events and :
 - (i) If $P(A) = 0.4$, $P(A \cup B) = 0.7$ then find $P(B)$.
 - (ii) if $P(A) = 0.5$, $P(A \cup B) = 0.7$ then find $P(B)$.
4. An urn contains 7 red and 4 blue balls. Two balls are drawn at random with replacement. Find the probability of getting : (a) 2 red balls, (b) 2 blue balls.
5. A bag contains 3 white and 5 black balls. Two balls are drawn at random without replacement. Determine the probability of getting the black balls.
6. A husband and wife appear in an interview for two vacancies in the same post. The probability of husband's selection is $1/7$ and that of wife's is $1/5$. Find the probability that (a) both get selected (b) only one of them get selected.
7. The probability of A hitting a target is $4/5$ and that of B is $2/3$. They both fire at the target. Find the probability that : (a) at least one of them will hit the target, (b) only one of them will hit the target.
8. A problem is given to 3 students whose chances of solving it are $1/3$, $1/5$ and $1/6$. What is the probability that (i) exactly two of them may solve it, (ii) at least two of them will solve it, (iii) problem will be solved.
9. Two bags contain 6 red and 4 black balls, 3 red and 3 black balls. One ball is drawn at random from one of the bags and it is found to be red. Find the probability that it was drawn from first bag.
10. Two bags contain 7 red and 2 black balls, 3 red and 6 black balls. One ball is drawn at random from one of the bags and it is found to be red. Find the probability that it was drawn from first bag.
11. Two bags contain 6 red and 8 black balls, 9 red and 7 black balls. One ball is drawn at random from one of the bags. Find the probability of drawing a black ball.
12. In a factory which manufactures bolts, machine A, B and C manufacture respectively 25%, 35% and 40% of the bolts. Of their outputs 5%, 4% and 2% are respectively defective bolts. A bolt is drawn at random from the product and is found to be defective. What is the probability that it is manufactured by the machine B.